arXiv:cond-mat/0211655v1 [cond-mat.str-el] 28 Nov 2002

# Exact solution for the two-site Kondo-Lattice Model - a limiting case for an insulator

T. Hickel,* J. Röseler, and W. Nolting
*Lehrstuhl für Festkörpertheorie, Institut für Physik, Humboldt-Universität, 10115 Berlin*
(Dated: February 1, 2008)

The Kondo-lattice model is well established as a method to describe an exchange coupling between single conduction electrons and localized magnetic moments. As a nontrivial exact result the zero-bandwidth limit (*atomic limit*) can be used to test approximations for this model. As soon as the translational symmetry is broken (for instance by sublattice structures) it is necessary to consider more than one lattice site. Therefore, we study as a starting point for generalizations the situation of a two-site cluster. An equation-of-motion approach is chosen to obtain the one-particle Green's function. In order to determine the spectral weights of its energy poles, we derive different possibilities for the calculation of the involved correlation functions. In this paper the analytical exact result for the situation of an insulator is presented. In a forthcoming article we generalize the calculations to arbitrary electron densities, keeping as the only constraint $S = 1/2$.

## I. INTRODUCTION

The idea of the Kondo lattice model goes back to Zener[1], one of the first scientists who tried to give a qualitative explanation of the physical behaviour of transition metals. His model is based on the existence of a local magnetic moment $\mathbf{S}_i$ per lattice site $i$ composed by electrons of partially filled electron shells. This assumption is well fulfilled in the 3d and 4f shells of transition metals and lanthanides respectively, since these shells are screened very effectively by outer filled electron shells. A strong Hund's rule coupling within the shells in connection with the notion of the Wigner-Eckart theorem leads to a well defined quantum number $S$ given by $\mathbf{S}_i^2 = \hbar^2 S(S+1)$. Due to exchange mechanisms there exists an effective coupling between the localized spins, which has the Heisenberg form

$$\mathcal{H}_{ff} = -\sum_{i,j} J_{ij}\, \mathbf{S}_i \cdot \mathbf{S}_j. \tag{1}$$

The electronic behaviour of these materials is determined by another class of electrons, namely the 5s and 5d/6s shells of the 3d- and 4f-systems, respectively. These electrons are itinerant and their uncorrelated propagation through the lattice is described by the Hamiltonian

$$\mathcal{H}_s = \sum_{i,j} \sum_{\sigma} T_{ij} c_{i\sigma}^{\dagger} c_{j\sigma}. \tag{2}$$

Using second quantization, the Fermi operator $c_{j\sigma}$ annihilates an electron with spin $\sigma$ at site $j$ whereas $c_{i\sigma}^{\dagger}$ creates one at site $i$. The hopping integrals $T_{ij}$ are connected by Fourier transformation to the single electron Bloch energies

$$T_{ij} = \frac{1}{N} \sum_{\mathbf{k}} \epsilon(\mathbf{k}) \mathrm{e}^{\mathrm{i}\mathbf{k}(\mathbf{R}_i - \mathbf{R}_j)}. \tag{3}$$

Zener pointed out that for the transition metals it is insufficient to handle the magnetic and the electronic partial system independently. Instead, he suggested an on-site interaction of the itinerant electron spin $\boldsymbol{\sigma}$ with the localized magnetic moments $\mathbf{S}_i$ of the form

$$\mathcal{H}_{sf} = -\frac{J}{\hbar} \sum_i \boldsymbol{\sigma}_i \cdot \mathbf{S}_i. \tag{4}$$

Due to the close relation to Kondo systems, the Hamiltonian $\mathcal{H}_s + \mathcal{H}_{sf}$ is often called *Kondo lattice model*[2]. Another common name is *s-f model* or *s-d model* emphasizing the class of described materials. In the case of manganites the same Hamiltonian is discussed in the limit $J\hbar \gg T_{ij}$ and called *double exchange model*, based on a special mechanism for the hopping of electrons in these insulators. Furthermore, one might note that the *periodic Anderson model* can also be mapped on a Hamiltonian of the form $\mathcal{H}_s + \mathcal{H}_{sf}$ in the limiting case of small charge fluctuations.

Sometimes, especially if higher electron concentrations are considered, it is reasonable to include correlations within the itinerant electron partial system. The simplest possibility is that of an on-site Coulomb repulsion

$$\mathcal{H}_{ss} = \frac{U}{2} \sum_i \sum_{\sigma} \hat{n}_{i\sigma} \hat{n}_{i-\sigma} = U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow}. \tag{5}$$

We are interested in the composed Hamiltonian

$$\mathcal{H} = \mathcal{H}_s + \mathcal{H}_{ss} + \mathcal{H}_{sf} + \mathcal{H}_{ff}, \tag{6}$$

which we shall call **correlated Kondo lattice model** (CKLM). By making certain assumptions for the parameters $T_{ij}, U, J_{ij}$ and $J$ one might return to somewhat more special situations.

The CKLM is not exactly soluble. One needs approximation schemes, which are often difficult to justify. Limiting cases are a powerful tool to weaken this shortcoming.

Till now only a few exact statements on the CKLM are known. One of them concerns the limiting case of a single excess electron in an otherwise empty conduction band, interacting with a ferromagnetically saturated system of localized spins ($T = 0\,\mathrm{K}$). Under these assumptions the electron may exist in a bound state, called magnetic polaron[3,4], which for $J < 0$ can be shown to be the ground state[5].

Another known and often used exact result for the CKLM is that of the zero-bandwidth limit[6]. Here one assumes the energy dispersion to be flat: $\epsilon(\mathbf{k}) \to T_0$, and is then able to calculate all possible one-particle excitation energies together with the corresponding spectral weights. Due to the relation (3) the assumption is equivalent to $T_{ij} \to \delta_{ij} T_0$, meaning that the *a priori* itinerant electrons described by $\mathcal{H}_s + \mathcal{H}_{ss}$ are not allowed to change the lattice site. For this reason the zero-bandwidth limit describes a one-site situation, sometimes called *atomic limit*.

With this paper we intend to improve the atomic limit result by going one step further towards the lattice. A two-site cluster possesses two main advantages: First, a non-zero intersite hopping integral allows a motion of electrons and leads to a finite energy dispersion. Compared to the flat energy band of the atomic limit, this is a qualitative improvement. Secondly, at least two lattice sites are necessary to provide a limiting case for an antiferromagnet. After the two-site CKLM has been solved exactly, we suppose to have a powerful tool to test approximation schemes for antiferromagnetic solutions of the full CKLM (6). Furthermore, we hope that the knowledge about properties and symmetries we gained when dealing with the two-site problem as well as the techniques we used to solve it are helpful tools to develop such approximate schemes.

Especially for the Hubbard model $\mathcal{H}_s + \mathcal{H}_{ss}$ there have already been some attempts to generalize exact cluster-results to the lattice[7,8,9,10] and to solve even four-site clusters exactly[11]. In this context one should also mention efforts to perform a cluster CPA for the Hubbard model.

For the CKLM a complete analytical solution for the one-particle Green's function is so far missing. Only Matlak *et al.*[12] have dealt with the eigenvalue problem of a two-site cluster for $S = 1/2$ and gave some results for certain correlation functions. With our work we tried to tackle the task of the exact solution for a two-site cluster more comprehensively. The present article is devoted to the situation of an insulator, being characterized by an empty conduction band. It is our intention to explain the employed mathematical techniques and to discuss intensively the difficulty of a connection between cluster and lattice. In a forthcoming article[13] we will generalize the calculations to arbitrary band occupations.

## II. CLUSTER MODEL

The Hamiltonian (6) possesses with $\mathcal{H}_s$ and $\mathcal{H}_{ff}$ two non-local terms. We mentioned before that a two-site cluster has two remarkable properties: it is sufficient for our intension to improve the atomic-limit result and it is still analytically manageable. These desired properties guided us to treat the non-local terms as follows:

The hopping integral is supposed to be

$$T_{ij} = \begin{cases} \tilde{T} & i \neq j \text{ and } i,j \text{ in same cluster,} \\ T_0 & i = j\,, \\ 0 & i,j \text{ in different clusters,} \end{cases}$$

leading to a situation as shown in figure 1.




FIG. 1: Formation of two-site clusters due to a restriction of the electron hopping. Different symbols for the lattice sites are chosen to adumbrate an ABAB-ordered antiferromagnet.

In our calculations, the Heisenberg term $\mathcal{H}_{ff}$ is excluded from the Hamiltonian. This results in a quasi separation of the spin system and the electronic system. The calculation of the dynamics of localized spins is placed back in favour of a description of the conduction band. However, a specification of the first influences the latter. This offers a chance to connect the two-site cluster to the lattice. A localized magnetic moment, calculated within the cluster, would give a vanishing value. If (anti-)ferromagnetic substances are the subject of investigation, one should instead specify the values for $\langle S_1^z \rangle$ and $\langle S_2^z \rangle$ to be finite. This is only possible if the Hamiltonian of the cluster itself does not include a Heisenberg interaction of the localized spins. This assumption about our model is also justified since in many papers the Kondo-lattice model is understood to consist only of $\mathcal{H}_s + \mathcal{H}_{sf}$ anyway[2]. Matlak *et al.*[12] have chosen another possibility by considering a mean-field approximated Heisenberg term as part of the Hamiltonian. Then one is confronted with even more difficult mathematical expressions than given here.

A simplification we make in this particular article regards the number of electrons in the conduction band $N \equiv \sum n_{i\sigma}$. As mentioned before, we assume $N = 0$ to be able to give a concise illustration of mathematical techniques and physical properties. As a consequence, all

expectation values which, like $\langle \hat{n}_{1\uparrow} \rangle$, $\langle S_2^- \hat{n}_{2\uparrow} c_{1\uparrow}^\dagger c_{1\downarrow} \rangle$, ..., possess in normal order an annihilation operator are zero. However, a treatment with Green's functions implies the existence of an additional "test electron", still giving rise to non-trivial results. Nevertheless, the term $\mathcal{H}_{ss}$ of the CKLM, which describes a Coulomb-interaction of *two* conduction electrons, is meaningless in the limit $N = 0$. Therefore, it can be omitted.

Altogether, our cluster model is described by the operator ( $S^\uparrow \equiv S^+, S^\downarrow \equiv S^-, z_\uparrow \equiv +1, z_\downarrow \equiv -1$ ):

$$\begin{aligned} \bar{\mathcal{H}} &= \tilde{T} \sum_{\sigma=\uparrow,\downarrow} \left( c_{1\sigma}^\dagger c_{2\sigma} + c_{2\sigma}^\dagger c_{1\sigma} \right) \qquad (7) \\ &+ \sum_{\alpha=1}^{2} \sum_{\sigma=\uparrow,\downarrow} \left\{ T_0 \hat{n}_{\alpha\sigma} - \frac{J}{2} \left( z_\sigma S_\alpha^z \hat{n}_{\alpha\sigma} + S_\alpha^\sigma c_{\alpha-\sigma}^\dagger c_{\alpha\sigma} \right) \right\}. \end{aligned}$$

Because it is a finite system, any physical quantity can in principle be calculated exactly. Without any restriction for the concentration of conduction electrons, the dimension of the Hilbert space would be $4^2 \cdot (2S+1)^2$. We present here results for $S = 1/2$. For this special case one can make use of the operator identity $\left(S_i^z\right)^2 \equiv \hbar^2 S^2 \mathbb{1}$, which allows to replace any product of two spin operators at the same site by a single one:

$$S_i^{\pm\sigma} S_i^{\mp\sigma} = \hbar^2 S \pm \hbar z_\sigma S_i^z, \qquad (8)$$

$$S_i^{\pm\sigma} S_i^z = -S_i^z S_i^{\pm\sigma} = \mp \hbar S z_\sigma S_i^{\pm\sigma}. \qquad (9)$$

Comparable relations do also exist[14] for $S > 1/2$, giving a hint that the qualitative structure of our results is also valid for other values of $S$.

Even so we are able to solve the eigenvalue problem for every electron concentration, it is our intension to handle the cluster with a many-particle approach. The effort to determine the complete single-particle excitation spectrum of the cluster is comparable for both techniques. Additionally, the calculation of the one-electron Green's function provides information on the dependence of the spectral weights not only on the model parameters but also on certain expectation values as the averaged $z$-component of the localized spins. However, the main reason for our choice is the wish to get a deeper understanding in the physics and possible approximations of the full CKLM, where a many-particle treatment is inevitable.

## III. SOLUTION FOR THE EQUATIONS OF MOTION

Due to the Heisenberg time dependence of the operators the retarded one-particle Green's function

$$\left\langle\!\left\langle \hat{A}(t); \hat{B}(t') \right\rangle\!\right\rangle := -\mathrm{i}\Theta(t-t') \left\langle \left[ \hat{A}(t), \hat{B}(t') \right]_+ \right\rangle \qquad (10)$$

has the following equation of motion

$$E \left\langle\!\left\langle \hat{A}; \hat{B} \right\rangle\!\right\rangle_E = \hbar \left\langle [\hat{A}, \hat{B}]_+ \right\rangle + \left\langle\!\left\langle [\hat{A}, \bar{\mathcal{H}}]_-; \hat{B} \right\rangle\!\right\rangle_E. \qquad (11)$$

Here, we used the Fourier-transformed notation

$$\left\langle\!\left\langle \hat{A}; \hat{B} \right\rangle\!\right\rangle_E := \int_{-\infty}^{\infty} \mathrm{d}(t-t') \left\langle\!\left\langle \hat{A}(t); \hat{B}(t') \right\rangle\!\right\rangle \exp\left[ \frac{\mathrm{i}}{\hbar} E(t-t') \right].$$

Motivated by the site symmetry, it is convenient to consider the following combinations of Green's functions:

$$\begin{aligned} \left\langle\!\left\langle \hat{C}_s \hat{D}_{\bar{s}} c_{s\sigma} \right\rangle\!\right\rangle^{(\epsilon)} &= \left\langle\!\left\langle \hat{C}_s \hat{D}_{\bar{s}} c_{s\sigma}; c_{s\sigma}^\dagger \right\rangle\!\right\rangle_E \\ &+ \epsilon \left\langle\!\left\langle \hat{C}_{\bar{s}} \hat{D}_s c_{\bar{s}\sigma}; c_{s\sigma}^\dagger \right\rangle\!\right\rangle_E. \qquad (12) \end{aligned}$$

Here, the site index $\bar{s}$ represents the opposite of site $s$ ($s = 1 \Rightarrow \bar{s} = 2; s = 2 \Rightarrow \bar{s} = 1$), $\hat{C}_s$ and $\hat{D}_{\bar{s}}$ are arbitrary products of spin operators at site $s$ and $\bar{s}$, respectively, $\epsilon = +1$ or $-1$.

We are especially interested in the one-electron Green's function $\left\langle\!\left\langle c_{i\sigma}; c_{j\sigma}^\dagger \right\rangle\!\right\rangle_E$ or $\langle\!\langle c_{s\sigma} \rangle\!\rangle^{(\epsilon)}$. The complete set of Green's functions generated by a repeated implementation of (11) is:

$$\begin{array}{lllll} G_1^{(\epsilon)} &=& \langle\!\langle c_{s\sigma} \rangle\!\rangle^{(\epsilon)} & K_1 &= \hbar, \\ G_2^{(\epsilon)} &=& \langle\!\langle S_s^z c_{s\sigma} \rangle\!\rangle^{(\epsilon)} & K_2 &= \hbar \langle S_s^z \rangle, \\ G_3^{(\epsilon)} &=& \langle\!\langle S_{\bar{s}}^z c_{s\sigma} \rangle\!\rangle^{(\epsilon)} & K_3 &= \hbar \langle S_{\bar{s}}^z \rangle, \\ G_4^{(\epsilon)} &=& \left\langle\!\left\langle S_s^{-\sigma} c_{s-\sigma} \right\rangle\!\right\rangle^{(\epsilon)} & K_4 &= 0, \\ G_5^{(\epsilon)} &=& \left\langle\!\left\langle S_{\bar{s}}^{-\sigma} c_{s-\sigma} \right\rangle\!\right\rangle^{(\epsilon)} & K_5 &= 0, \\ G_6^{(\epsilon)} &=& \langle\!\langle S_s^z S_{\bar{s}}^z c_{s\sigma} \rangle\!\rangle^{(\epsilon)} & K_6 &= \hbar \langle S_s^z S_{\bar{s}}^z \rangle, \\ G_7^{(\epsilon)} &=& \left\langle\!\left\langle S_{\bar{s}}^z S_s^{-\sigma} c_{s-\sigma} \right\rangle\!\right\rangle^{(\epsilon)} & K_7 &= 0, \\ G_8^{(\epsilon)} &=& \left\langle\!\left\langle S_s^z S_{\bar{s}}^{-\sigma} c_{s-\sigma} \right\rangle\!\right\rangle^{(\epsilon)} & K_8 &= 0, \\ G_9^{(\epsilon)} &=& \left\langle\!\left\langle S_{\bar{s}}^{-\sigma} S_s^\sigma c_{s\sigma} \right\rangle\!\right\rangle^{(\epsilon)} & K_9 &= \hbar \langle S_{\bar{s}}^{-\sigma} S_s^\sigma \rangle, \\ G_{10}^{(\epsilon)} &=& \langle\!\langle S_s^{-\sigma} S_{\bar{s}}^\sigma c_{s\sigma} \rangle\!\rangle^{(\epsilon)} & K_{10} &= \hbar \langle S_s^{-\sigma} S_{\bar{s}}^\sigma \rangle. \end{array} \qquad (13)$$

In the right column the corresponding inhomogeneities in the equation of motion (11) to each of the Green's functions is given. Emerging Green's functions with another combination of spin operators or a higher number of electron operators vanish identically because of the constraints $S = 1/2$ and $N = 0$.

It is a valuable property of the combination (12) that the equations of motion of the $G^{(+)}$ and the $G^{(-)}$ Green's functions do not mix. Moreover, if one writes (11) in the form

$$\sum_{j=1}^{10} m_{ij} G_j^{(\epsilon)} = K_i, \qquad (14)$$

one gets a matrix $M = (m_{ij})_{i,j}$ which contains $\epsilon$ only as a pre-factor of $\tilde{T}$. If the matrix equation (14) is solved for $\epsilon = +1$, one obtains the expression for $G_i^{(-)}$ simply by replacing in $G_i^{(+)}$ the parameter $\tilde{T}$ by $-\tilde{T}$. We therefore omit $\epsilon$ in the forthcoming equations.

By a couple of unitary transformations $M$ can be reduced further to blocks of a size not greater than $3\times 3$. It is favourable to introduce the combined Green's functions

$$\begin{aligned}
H_0^{(\pm)} &= \tfrac{\hbar}{2}\langle\langle c_{s\sigma}\rangle\rangle \pm z_\sigma \langle\langle S_s^z c_{s\sigma}\rangle\rangle \\
&\qquad = \tfrac{1}{\hbar}\langle\langle S_s^{\pm\sigma} S_s^{\mp\sigma} c_{s\sigma}\rangle\rangle, \\
H_1^{(\pm)} &= \tfrac{\hbar}{2}\langle\langle S_{\bar s}^z c_{s\sigma}\rangle\rangle \pm z_\sigma \langle\langle S_s^z S_{\bar s}^z c_{s\sigma}\rangle\rangle \\
&\qquad = \tfrac{1}{\hbar}\langle\langle S_s^{\pm\sigma} S_s^{\mp\sigma} S_{\bar s}^z c_{s\sigma}\rangle\rangle, \\
H_2^{(\pm)} &= \tfrac{\hbar}{2} H_0^{(\pm)} \pm z_\sigma H_1^{(\pm)} \\
&\qquad = \tfrac{1}{\hbar^2}\langle\langle S_s^{\pm\sigma} S_s^{\mp\sigma} S_{\bar s}^{\pm\sigma} S_{\bar s}^{\mp\sigma} c_{s\sigma}\rangle\rangle, \\
H_3^{(\pm)} &= \tfrac{\hbar}{2} H_0^{(\mp)} \pm z_\sigma H_1^{(\mp)} \\
&\qquad = \tfrac{1}{\hbar^2}\langle\langle S_s^{\mp\sigma} S_s^{\pm\sigma} S_{\bar s}^{\pm\sigma} S_{\bar s}^{\mp\sigma} c_{s\sigma}\rangle\rangle, \\
H_4^{(\pm)} &= \tfrac{\hbar}{2}\langle\langle S_s^{-\sigma} c_{s-\sigma}\rangle\rangle \pm z_\sigma \langle\langle S_{\bar s}^z S_s^{-\sigma} c_{s-\sigma}\rangle\rangle \\
&\qquad = \tfrac{1}{\hbar}\langle\langle S_{\bar s}^{\pm\sigma} S_{\bar s}^{\mp\sigma} S_s^{-\sigma} c_{s\sigma}\rangle\rangle, \\
H_5^{(\pm)} &= \tfrac{\hbar}{2}\langle\langle S_{\bar s}^{-\sigma} c_{s-\sigma}\rangle\rangle \pm z_\sigma \langle\langle S_s^z S_{\bar s}^{-\sigma} c_{s-\sigma}\rangle\rangle \\
&\qquad = \tfrac{1}{\hbar}\langle\langle S_s^{\pm\sigma} S_s^{\mp\sigma} S_{\bar s}^{-\sigma} c_{s\sigma}\rangle\rangle .
\end{aligned} \tag{15}$$

In another step the combinations $H_3^{(-)} \pm G_{10}, H_3^{(+)} \pm G_9$ and $H_4^{(+)} \pm H_5^{(+)}$ are used, what finishes the reduction.

The remaining $3\times 3$ blocks have the following matrix-structure:

$$M_0^{(\mu_1\mu_2\mu_3\mu_4)}(\hat E) = \begin{pmatrix} \hat E + 2\mu_4 a & \mu_1\tilde T & 0 \\ \mu_1\tilde T & \hat E & \mu_2 2a \\ 0 & \mu_2 2a & \hat E + \mu_3\tilde T \end{pmatrix} \tag{16}$$

Here, we used the abbreviation $a = \frac{\hbar}{2}\frac{J}{2}$ and $\mu_1,\ldots,\mu_4$ are sign parameters. The eigenvalues of this matrix are

$$\begin{aligned}
\hat E_{01}^{(\mu_3\mu_4)} &= \mu_3\tilde T + 2\mu_4 a, \\
\hat E_{02}^{(\mu_3\mu_4)} &= -\sqrt{4a^2 - 2\mu_3\mu_4 a\tilde T + \tilde T^2}, \\
\hat E_{03}^{(\mu_3\mu_4)} &= +\sqrt{4a^2 - 2\mu_3\mu_4 a\tilde T + \tilde T^2}.
\end{aligned}$$

For the eigenvectors one gets

$$v_{01}^{(\mu_1\mu_2\mu_3\mu_4)} = \frac{1}{\sqrt 3}\begin{pmatrix} \mu_1\mu_3 \\ 1 \\ \mu_2\mu_4 \end{pmatrix} \tag{17}$$

and for $k = 2, 3$

$$v_{0k}^{(\mu_1\mu_2\mu_3\mu_4)} = p_k \begin{pmatrix} \mu_1(\tilde T - 2\mu_3\mu_4 a - \mu_3\hat E_{0k}) \\ -\mu_3\tilde T + \hat E_{0k} \\ 2\mu_2 a \end{pmatrix} \tag{18}$$

$$\text{with}\quad p_k = \frac{1}{2a}\sqrt{\frac{\hat E_{0k} + \mu_3\tilde T - \mu_4 a}{3\hat E_{0k}}}. \tag{19}$$

We denote by $v_{0k}[i]$ the $i$th component of $v_{0k}$ and define a matrix $V$ by its components $V_0[ik] = v_{0k}[i]$. Basic algebra shows that $M_0 = V_0\cdot D_0\cdot V_0^{-1}$ with $D_0 = \mathrm{diag}(\hat E + \hat E_{0k})$. Using these expressions one can give the components of the inverse of $M_0$ in a form which is characterized by linear energy poles:

$$M_0^{-1}[ij] = \sum_{k=1}^{3} v_{0k}[i]v_{0k}[j]\frac{1}{\hat E + \hat E_{0k}}. \tag{20}$$

This form is particularly suitable for the calculation of Green's functions.

The $10\times 10$ matrix equation (14) can now be written in the form

$$\begin{aligned}
M_0^{(-+-+)}(E - T_0 - a)\begin{pmatrix} H_5^{(-)} \\ H_4^{(-)} \\ H_2^{(-)} \end{pmatrix} &= \begin{pmatrix} R_1 \\ R_2 \\ R_3 \end{pmatrix}, \\
M_0^{(-+-+)}(E - T_0 - a)\begin{pmatrix} H_3^{(-)} + G_{10} \\ H_3^{(+)} + G_9 \\ H_4^{(+)} + H_5^{(+)} \end{pmatrix} &= \begin{pmatrix} R_4 + R_7 \\ R_5 + R_8 \\ R_6 + R_9 \end{pmatrix}, \\
M_0^{(-+++)}(E - T_0 - a)\begin{pmatrix} H_3^{(-)} - G_{10} \\ H_3^{(+)} - G_9 \\ H_4^{(+)} - H_5^{(+)} \end{pmatrix} &= \begin{pmatrix} R_4 - R_7 \\ R_5 - R_8 \\ R_6 - R_9 \end{pmatrix}
\end{aligned}$$

and solved simply by applying equation (20). The emerging inhomogeneities $R_1,\ldots,R_9$ are constructed out of $K_1,\ldots,K_{10}$, obeying the same rules of combination as for the Green's functions. The single missing equation has an even simpler structure. One can write down immediately the solution for $H_2^{(+)}$:

$$\begin{aligned}
H_2^{(+)} &= \frac{R_0}{E - T_0 - \tilde T + a} \\
&= \frac{\frac{\hbar}{2}\left(\frac{1}{2}\hbar^2 + \hbar z_\sigma\langle S_1^z\rangle\right) + \left(\frac{1}{2}\hbar^2 z_\sigma\langle S_2^z\rangle + \hbar\langle S_1^z S_2^z\rangle\right)}{E - T_0 - \tilde T + a}
\end{aligned} \tag{21}$$

To obtain an expression for the one-particle Green's function we are looking for, one has to take a sum of the combined functions $H_2^{(\pm)},\ldots H_5^{(\pm)}$. Hence,

$$\begin{aligned}
\langle\langle c_{s\sigma}\rangle\rangle = G_1 &= \frac{1}{\hbar^2}\left(H_2^{(+)} + H_3^{(+)} + H_2^{(-)} + H_3^{(-)}\right) \\
&= \frac{1}{\hbar^2}\left(H_2^{(+)} + H_2^{(-)}\right) \\
&+ \frac{1}{2\hbar^2}\left(\left(H_3^{(-)} + G_{10}\right) + \left(H_3^{(+)} + G_9\right)\right) \\
&+ \frac{1}{2\hbar^2}\left(\left(H_3^{(-)} - G_{10}\right) + \left(H_3^{(+)} - G_9\right)\right).
\end{aligned} \tag{22}$$

We now remind the fact stated after equation (14) that this solution can be obtained equally well for $\epsilon = +1$ as for $\epsilon = -1$. If we add these two Green's functions

$$\left\langle\langle c_{s\sigma}; c_{s\sigma}^\dagger\right\rangle\rangle_E = \frac{1}{2}\left(\langle\langle c_{s\sigma}\rangle\rangle^{(+)} + \langle\langle c_{s\sigma}\rangle\rangle^{(-)}\right), \tag{23}$$

we effectively reverse the combination (12), which was a consequence of site symmetry.

Therefore, the following result for the one-particle Green's function is obtained:

$$\begin{aligned}
&\left\langle\langle c_{s\sigma}; c_{s\sigma}^\dagger\right\rangle\rangle_E \\
&= \left(\frac{\hbar}{E - \varepsilon_1} + \frac{\hbar}{E - \varepsilon_2}\right)\cdot \\
&\quad\cdot\frac{1}{6\hbar^2}\left(\frac{3}{2}\hbar^2 + z_\sigma\hbar\langle S_x^z\rangle + z_\sigma\hbar\langle S_y^z\rangle\right.
\end{aligned}$$

$$
\begin{aligned}
&+2\langle S_x^z S_y^z\rangle + \langle S_y^{-\eta} S_x^{\eta}\rangle + \langle S_x^{-\eta} S_y^{\eta}\rangle\big) \\
- &\left(\frac{\hbar/\hat{E}_{02}^{(-+)}}{E-\varepsilon_3} + \frac{\hbar/\hat{E}_{03}^{(-+)}}{E-\varepsilon_5} + \frac{\hbar/\hat{E}_{02}^{(++)}}{E-\varepsilon_4} + \frac{\hbar/\hat{E}_{03}^{(++)}}{E-\varepsilon_6}\right)\cdot \\
&\cdot\frac{\hbar\frac{J}{2}}{12\hbar^2}\big(-2z_\sigma\hbar\langle S_x^z\rangle + z_\sigma\hbar\langle S_y^z\rangle + 2\langle S_x^z S_y^z\rangle \\
&+\langle S_y^{-\eta} S_x^{\eta}\rangle + \langle S_x^{-\eta} S_y^{\eta}\rangle\big) \\
+ &\left(\frac{\hbar}{E-\varepsilon_3} + \frac{\hbar}{E-\varepsilon_5} + \frac{\hbar}{E-\varepsilon_4} + \frac{\hbar}{E-\varepsilon_6}\right)\cdot \\
&\cdot\frac{1}{12\hbar^2}\left(\frac{3}{2}\hbar^2 - z_\sigma\hbar\langle S_x^z\rangle - z_\sigma\hbar\langle S_y^z\rangle\right. \\
&\left.-2\langle S_x^z S_y^z\rangle - \langle S_y^{-\eta} S_x^{\eta}\rangle - \langle S_x^{-\eta} S_y^{\eta}\rangle\right) \\
- &\left(\frac{\hbar/\hat{E}_{02}^{(-+)}}{E-\varepsilon_3} + \frac{\hbar/\hat{E}_{03}^{(-+)}}{E-\varepsilon_5} - \frac{\hbar/\hat{E}_{02}^{(++)}}{E-\varepsilon_4} - \frac{\hbar/\hat{E}_{03}^{(++)}}{E-\varepsilon_6}\right)\cdot \\
&\cdot\frac{\tilde{T}}{12\hbar^2}\big(-z_\sigma\hbar\langle S_x^z\rangle - z_\sigma\hbar\langle S_y^z\rangle + 4\langle S_x^z S_y^z\rangle \\
&+2\langle S_y^{-\eta} S_x^{\eta}\rangle + 2\langle S_x^{-\eta} S_y^{\eta}\rangle\big)
\end{aligned}
\tag{24}
$$

It has the six energy poles

$$
\begin{aligned}
\varepsilon_1 &= T_0 + a - \hat{E}_{01}^{(-+)} = T_0 - \tfrac{\hbar}{2}\tfrac{J}{2} + \tilde{T}, \\
\varepsilon_2 &= T_0 + a - \hat{E}_{01}^{(++)} = T_0 - \tfrac{\hbar}{2}\tfrac{J}{2} - \tilde{T}, \\
\varepsilon_3 &= T_0 + a - \hat{E}_{02}^{(-+)} = T_0 + \tfrac{\hbar}{2}\tfrac{J}{2} + \sqrt{\left(\hbar\tfrac{J}{2}\right)^2 + \hbar\tfrac{J}{2}\tilde{T} + \tilde{T}^2}, \\
\varepsilon_4 &= T_0 + a - \hat{E}_{02}^{(++)} = T_0 + \tfrac{\hbar}{2}\tfrac{J}{2} + \sqrt{\left(\hbar\tfrac{J}{2}\right)^2 - \hbar\tfrac{J}{2}\tilde{T} + \tilde{T}^2}, \\
\varepsilon_5 &= T_0 + a - \hat{E}_{03}^{(-+)} = T_0 + \tfrac{\hbar}{2}\tfrac{J}{2} - \sqrt{\left(\hbar\tfrac{J}{2}\right)^2 + \hbar\tfrac{J}{2}\tilde{T} + \tilde{T}^2}, \\
\varepsilon_6 &= T_0 + a - \hat{E}_{03}^{(++)} = T_0 + \tfrac{\hbar}{2}\tfrac{J}{2} - \sqrt{\left(\hbar\tfrac{J}{2}\right)^2 - \hbar\tfrac{J}{2}\tilde{T} + \tilde{T}^2},
\end{aligned}
$$

shown in figure 2.




FIG. 2: The hybridization $\tilde{T}$ causes a splitting of the one-electron energy levels valid for a single site and a grouping of the excitation energies. For $\tilde{T} = 0.5\,\mathrm{eV}$ and $J\hbar = 0.3\,\mathrm{eV}, T_0 = 0\,\mathrm{eV}$ the arrangement of the energy poles $E = \varepsilon_2, \varepsilon_5, \varepsilon_6, \varepsilon_1, \varepsilon_4, \varepsilon_3$ (starting from the bottom) is shown.

## IV. TREATMENT OF THE CORRELATION FUNCTIONS

The result (24) provides an expression for the one-particle Green's function. It is characterized by six linear energy poles in the denominators. They are identical with the excitation energies possible if a single excess electron is placed in the otherwise empty conduction band. The numerators give the spectral weights corresponding to these excitations. However, in contrast to the model parameters, $\tilde{T}, T_0, J$, which are assumed to be known, we have till now no information about the spin correlation functions $\langle S_1^z\rangle, \langle S_2^z\rangle, \langle S_1^z S_2^z\rangle, \langle S_1^+ S_2^-\rangle, \langle S_1^- S_2^+\rangle$, leaving the spectral weights undetermined.

In the next two subsections we will introduce and discuss two possible concepts to obtain results for these correlation functions. They are especially important for the treatment of a situation with $N > 0$.

### A. Remaining in the cluster

As a result of the calculations in section III we do not only get an expression for the Green's function $G_1^{(\epsilon)}$ and therefore for $\langle\langle c_{s\sigma}; c_{s\sigma}^\dagger\rangle\rangle_E$. On the contrary, we are without further effort able to write down similar expressions for all the other participating Green's functions $G_2^{(\epsilon)}, \ldots, G_{10}^{(\epsilon)}$. One just has to sum the combined Green's functions $H_2^{(\pm)}, \ldots, H_5^{(\pm)}$ another way as in (22).

It is one possible concept to apply the well-known spectral theorem to each of the determined Green's functions. If $G_1^{(\epsilon)}, G_2^{(\epsilon)}, \ldots$ or $G_{10}^{(\epsilon)}$ is of the form

$$
\left\langle\left\langle \hat{A}; \hat{B}\right\rangle\right\rangle_{E+\mathrm{i}0^+} = \sum_{i=1}^{p} \frac{\hbar\,\alpha_i}{E - E_i + \mathrm{i}0^+}, \tag{25}
$$

then one can derive an expression for a corresponding correlation function

$$
\left\langle \hat{B}\hat{A}\right\rangle = \sum_{i=1}^{p} \alpha_i\, f_-(E_i) \quad = \sum_{i=1}^{p} \frac{\alpha_i}{\mathrm{e}^{\beta(E_i-\mu)} + 1}, \tag{26}
$$

with the help of this theorem ($\beta = (k_\mathrm{B}T)^{-1}$). One gets a set of ten equations, which are linear in the unknown correlation functions. To solve this systems means that only properties of the cluster determine the correlations functions.

However, in our special situation of $N = 0$ each of the expectation values at the left hand side of (26) is supposed to vanish, because they all include an electron density operator. Then the solution of the set of equations becomes trivial. One has to choose $\mu \ll \varepsilon_k \forall k$, to fulfill

the requirements for the electron density. This leaves the set of spin correlation functions again undefined.

Hence, if $N = 0$, the application of the spectral theorem to the determined (that means electronic) set of Green's functions fails to provide any information about spin correlation functions. Instead other Green's functions can be used. The equations of motion for $\langle\langle S_1^+; S_1^- \rangle\rangle_E$, $\langle\langle S_2^+; S_2^- \rangle\rangle_E$ and $\langle\langle S_2^z S_1^+; S_1^- \rangle\rangle_E$ are particularly simple in the case $N = 0$. If the spectral theorem is applied to its results, one obtains the information that $\langle S_1^z \rangle, \langle S_2^z \rangle$ and $\langle S_1^z S_2^z \rangle$ are zero, respectively. However, even when using $\langle\langle S_1^+; S_2^- \rangle\rangle_E$ or $\langle\langle S_2^+; S_1^- \rangle\rangle_E$ it is not possible to get any expression for $\langle S_1^+ S_2^- \rangle$ and $\langle S_1^- S_2^+ \rangle$. For these correlation functions the expectation value has to be determined directly by calculating the trace of the canonical ensemble with the four possible eigenstates of $\bar{\mathcal{H}}$ for $N = 0$. The result is again zero. For an insulator it is not surprising that all spin correlation functions vanish if they are solely determined within the cluster.

### B. Connection to the lattice

To create a connection of the two-site cluster to the lattice, the spin correlation functions need to be specified outside the cluster. It is the advantage of an insulator that this can be done without coming into conflict with the spectral theorem applied to the cluster.

As explained in section II we have quasi separated the electronic system and the spin system by excluding $\mathcal{H}_{ff}$ from the Hamiltonian. As a different problem, we can now consider a situation of two localized spins mutually interacting via $\mathcal{H}_{ff}$ with $J_{ij} = J_{\mathrm{H}}$ and embedded in the lattice via an additional term in the Hamiltonian of the form

$$\mathcal{H}_f = -b\,(S_1^z + \eta S_2^z)\,. \tag{27}$$

Simultaneous to the choice of $J_{\mathrm{H}}$, the sign parameter $\eta$ allows to distinguish between ferromagnetic ($\eta = +1$) and antiferromagnet ($\eta = -1$) configurations. In the first case $b = B \cdot \frac{g_J \mu_{\mathrm{B}}}{\hbar}$ is as a molecular or Weiss field, in the latter case it describes the crystal anisotropy. The complete set of Green's functions for this problem is derived in appendix A. As a result the expressions (A13)-(A16) and (A21)-(A24) for a set of correlation functions are achieved.

This set is identical with the correlation functions we need for the calculation of the electronic part described by $\bar{\mathcal{H}}$. Determined in this way and used in (24), we are sure that not only $\langle S_1^z \rangle$ and $\langle S_2^z \rangle$ but also the two-site correlation functions $\langle S_1^z S_2^z \rangle, \langle S_1^+ S_2^- \rangle$ and $\langle S_1^- S_2^+ \rangle$ are described *consistently* by two experimentally accessible parameters $J_{\mathrm{H}}$ and $B$. We created an intelligible connection to the lattice.

Only after these considerations we are able to specify the behaviour of the electrons in a two-site (anti-)ferromagnetic Kondo lattice. If we assume a ferromagnet ($\langle S_1^z \rangle = \langle S_2^z \rangle$), we obtain a dependence of the spectral weights belonging to the six energy poles of the single-electron Green's function on the magnetization as shown in figure 3. The magnetic field $B$ is uniquely used to create a ferromagnetic order of the system of localized spins and does not affect the conduction electrons directly. To obtain a saturation of about 95% one needs extremely large exchange fields of about $B = 400\,\mathrm{T}$, as can also be estimated if the tendency to disorder, $k_{\mathrm{B}}T$, and to order, $\mu_{\mathrm{B}}B$, are set equal. One can clearly see that for these fields the energy poles $\varepsilon_2$ and $\varepsilon_1$ become dominant. This is not surprising, since the configuration where the excess electron and the two localized spins have all parallel spin has exactly this excitation energy. The observation that for $B = 0\,\mathrm{T}$ these two poles still possess the highest spectral weight is connected to the fact that additionally some one-electron eigenstates of the Hamiltonian $\bar{\mathcal{H}}$ with the $z$-component of the total spin $\pm(2S - 1/2)$ have the same energy. The symmetry of lattice sites 1 and 2 nicely enters the figure, too: It is the reason for similar structures in the groups around $E = -\tilde{T} = -0.5\,\mathrm{eV}$ and around $E = \tilde{T} = 0.5\,\mathrm{eV}$.




FIG. 3: For the energy poles of the single-electron Green's function, the dependence of the corresponding spectral weights on a magnetization $\langle S_1^z \rangle = \langle S_2^z \rangle$ parallel to the spin of the test electron $\sigma$ is given (left). The latter is adjusted by an assumed molecular field $B$ (right). The fixed model parameters are: $J_{\mathrm{H}}\hbar^2 = 10^{-3}\,\mathrm{eV}$, $J\hbar = 0.3\,\mathrm{eV}$, $\tilde{T} = 0.5\,\mathrm{eV}$ and $g_J = 1, T_0 = 0\,\mathrm{eV}$. The temperature is $80\,\mathrm{K}$.

As one can see in figure 4 the situation is quite different if the spin of the test electron is antiparallel to the alignment of the localized spins, induced by the applied magnetic field. Here, the energy poles $\varepsilon_3$ and $\varepsilon_6$ get maximum spectral weight when the field is increased. However, the site symmetry is not affected.

If an antiferromagnetic configuration of the localized spins ($\langle S_1^z \rangle = -\langle S_2^z \rangle$) is considered, there is again no change in the positions of the energy poles (figure 2). Of course this does not apply to their spectral weights. Comparing figure 5 to figure 3, one notices already for zero magnetization ($\langle S_1^z \rangle = 0 \Leftrightarrow B = 0$) a different distribution. The opposite sign for $J_{\mathrm{H}}$ affects the non-zero correlation functions $\langle S_1^z S_2^z \rangle$ and $\langle S_1^+ S_2^- \rangle$. As concerns the dependence on $B$, primarily the behaviour of the en-




FIG. 4: Same as figure 3 with the only difference that the spin of the test electron $\sigma$ is antiparallel to the direction of the magnetization.




FIG. 5: Same as figure 3, but with $\langle S_1^z\rangle = -\langle S_2^z\rangle$ and $J_{\mathrm{H}}\hbar^2 = -10^{-3}\,\mathrm{eV}$. $B$ is the magnetic anisotropy.

ergy poles $\varepsilon_5$ and $\varepsilon_4$ has changed. Their spectral weight does not tend to zero (ferromagnet), but rather increases with $B$. In general the antiferromagnetic configuration leads to smaller changes of the spectral weights, since the average $\frac{1}{2}\left(\langle S_1^z\rangle + \langle S_2^z\rangle\right) \equiv 0$ remains constant.

The term $\mathcal{H}_f$, which enters the determination of the correlation functions, leads for $\eta = -1$ to a break-down of the site symmetry. Therefore, the two groups of energy poles in figure 5 show a slightly different dependence on $B$. One has to distinguish between $\left\langle\!\left\langle c_{1\uparrow}; c_{1\uparrow}^\dagger \right\rangle\!\right\rangle_E \equiv \left\langle\!\left\langle c_{2\downarrow}; c_{2\downarrow}^\dagger \right\rangle\!\right\rangle_E$ and $\left\langle\!\left\langle c_{1\downarrow}; c_{1\downarrow}^\dagger \right\rangle\!\right\rangle_E \equiv \left\langle\!\left\langle c_{2\uparrow}; c_{2\uparrow}^\dagger \right\rangle\!\right\rangle_E$. The latter situation has been presented in figure 6. The dependence on the spin direction of the electron is of course much smaller than for a ferromagnetic configuration. In fact, apart from an interchange of the two groups of energy poles the distribution of the spectral weights shown in figure 6 is almost the same as in figure 5. The formation of these two groups is a result of hybridization of the




FIG. 6: Same as figure 5, but the Green's function is evaluated at the site which has a localized spin antiparallel to the electron spin.

energy levels of the two single sites, as shown in figure 2. The fact that the spectral weights of the energies in the lower group get the values of the higher group (and vice versa) if the spin direction of the test electron is changed, is again related to site symmetry.

Of course, all these features can be explained by a detailed analysis of the eigenvalue problem. However, the Hilbert spaces that need to be considered are relatively complex. For instance the subspace corresponding to a $z$-component of the total spin of $1/2$ is 6-dimensional, and hence every eigenvector is a linear combination of 6 different spin configurations. Therefore, the possible analysis is too lengthy to be presented here.

## V. SUMMARY

We have shown that the Hamiltonian $\bar{\mathcal{H}}$ can be treated such that an analytic expression for the one-particle Green's function as given in (24) is obtained. In order to do that we formulated the complete set of equations of motion as a single matrix equation, made use of the symmetries of the problem and combined Green's functions to reduce the matrix to blocks of size $3\times 3$, which can be solved. In this article we studied the case of an insulator ($N = 0$), but a generalization of these techniques to the situation in a metal is possible even so calculations are much less concise there.[13]

Furthermore, the connection to a lattice has been discussed. We propose to use the correlation functions in the spectral weights of the Green's functions for this purpose. A helpful tool is a two-site Heisenberg model which has been separated before from the Hamiltonian (6) and calculated in appendix A. By adding a term $\mathcal{H}_f$ we can simulate ferromagnetic and antiferromagnetic lattices. This

assumption about the system of localized spins results in certain features of the electronic partial system, visualized exemplary by the spectral weights given in figure 3 to 6. The site symmetry has a major influence on its distribution.

**Acknowledgments**

One of us (T. H.) gratefully acknowledges the financial support of the *Studienstiftung des deutschen Volkes*. This work also benefitted from the financial support of the *Sonderforschungsbereich SFB 290* of the Deutsche Forschungsgemeinschaft.

## APPENDIX A: THE TWO-SPIN PROBLEM

It is the purpose of this appendix to provide a consistent set of spin correlation functions as input for the two-site CKLM. This is done by discussing the Heisenberg-type Hamiltonian

$$\mathcal{H}_{\mathrm{A}} = -J_{\mathrm{H}}\left(S_1^+S_2^- + S_1^-S_2^+ + 2S_1^zS_2^z\right) - b\left(S_1^z + \eta S_2^z\right). \tag{A1}$$

To obtain correlation functions we use the spectral theorem, which needs certain Green's functions as an input. Within an equation-of-motion (11) approach these are:

$$\begin{array}{rclcrcl} G_{11} &=& \left\langle\!\left\langle S_1^-; S_1^+\right\rangle\!\right\rangle_E & & K_{11} &=& -2\hbar^2\left\langle S_1^z\right\rangle, \\ G_{21} &=& \left\langle\!\left\langle S_2^-; S_1^+\right\rangle\!\right\rangle_E & & K_{21} &=& 0, \\ \Gamma_{12} &=& \left\langle\!\left\langle S_1^zS_2^-; S_1^+\right\rangle\!\right\rangle_E & & L_{12} &=& \hbar^2\left\langle S_1^+S_2^-\right\rangle, \\ \Gamma_{21} &=& \left\langle\!\left\langle S_2^zS_1^-; S_1^+\right\rangle\!\right\rangle_E & & L_{21} &=& -2\hbar^2\left\langle S_1^zS_2^z\right\rangle. \end{array} \tag{A2}$$

On the right hand site we give the corresponding inhomogeneities which appear in the equations of motion:

$$(E + b\hbar)\, G_{11} = K_{11} + 2J_{\mathrm{H}}\hbar\left(\Gamma_{12} - \Gamma_{21}\right) \tag{A3}$$

$$(E + \eta b\hbar)\, G_{21} = K_{21} - 2J_{\mathrm{H}}\hbar\left(\Gamma_{12} - \Gamma_{21}\right) \tag{A4}$$

$$(E + \eta b\hbar)\, \Gamma_{12} = L_{12} + J_{\mathrm{H}}\frac{\hbar^3}{2}\left(G_{11} - G_{21}\right) \tag{A5}$$

$$(E + b\hbar)\, \Gamma_{21} = L_{21} - J_{\mathrm{H}}\frac{\hbar^3}{2}\left(G_{11} - G_{21}\right) \tag{A6}$$

Following the rules identical to what has been used in (15) the simplification

$$\begin{pmatrix} E + b\hbar \pm J_{\mathrm{H}}\hbar^2 & \mp J_{\mathrm{H}}\hbar^2 \\ \mp J_{\mathrm{H}}\hbar^2 & E + \eta b\hbar \pm J_{\mathrm{H}}\hbar^2 \end{pmatrix} \begin{pmatrix} \frac{\hbar}{2}G_{11} \pm \Gamma_{21} \\ \frac{\hbar}{2}G_{21} \pm \Gamma_{12} \end{pmatrix} = \begin{pmatrix} \frac{\hbar}{2}K_{11} \pm L_{21} \\ \frac{\hbar}{2}K_{21} \pm L_{12} \end{pmatrix} \tag{A7}$$

is possible. The determinant of the emerging matrix, $D^{(\pm)}$, has the form

$$\begin{aligned} \frac{1}{D^{(\pm)}} &= \frac{1}{2\sqrt{\cdots}}\left(\frac{1}{E \pm J_{\mathrm{H}}\hbar^2 + \frac{\hbar}{2}(1+\eta)b - \sqrt{\cdots}} - \frac{1}{E \pm J_{\mathrm{H}}\hbar^2 + \frac{\hbar}{2}(1+\eta)b + \sqrt{\cdots}}\right) \\ \text{with } \sqrt{\cdots} &= \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + \frac{1}{2}(1-\eta)b^2\hbar^2}. \end{aligned} \tag{A8}$$

By taking the inverse of the matrix in (A7) and after performing a partial fraction expansion one therefore obtains the combined Green's functions

$$\tfrac{\hbar}{2}G_{11} \pm \Gamma_{21} = -\hbar^3\left\langle S_1^z\right\rangle\left(\frac{(1-\eta)\frac{\hbar b}{4\sqrt{\cdots}} + \frac{1}{2}}{E - E_1^{(\pm)}} - \frac{(1-\eta)\frac{\hbar b}{4\sqrt{\cdots}} - \frac{1}{2}}{E - E_2^{(\pm)}}\right)$$

$$
\begin{aligned}
&\mp\ 2\hbar^2 \left\langle S_1^z S_2^z \right\rangle \left( \frac{(1-\eta)\frac{\hbar b}{4\sqrt{\cdots}} + \frac{1}{2}}{E - E_1^{(\pm)}} - \frac{(1-\eta)\frac{\hbar b}{4\sqrt{\cdots}} - \frac{1}{2}}{E - E_2^{(\pm)}} \right) \\
&-\ \left\langle S_1^+ S_2^- \right\rangle \frac{J_{\mathrm{H}}\hbar^4}{2\sqrt{\cdots}} \left( \frac{1}{E - E_1^{(\pm)}} - \frac{1}{E - E_2^{(\pm)}} \right),
\end{aligned}
$$

$$
\begin{aligned}
\tfrac{\hbar}{2} G_{21} \pm \Gamma_{12} \ =\ & \pm \left\langle S_1^z \right\rangle \frac{J_{\mathrm{H}}\hbar^5}{2\sqrt{\cdots}} \left( \frac{1}{E - E_1^{(\pm)}} - \frac{1}{E - E_2^{(\pm)}} \right) \\
&+\ \left\langle S_1^z S_2^z \right\rangle \frac{J_{\mathrm{H}}\hbar^4}{\sqrt{\cdots}} \left( \frac{1}{E - E_1^{(\pm)}} - \frac{1}{E - E_2^{(\pm)}} \right) \\
&\pm\ \hbar^2 \left\langle S_1^+ S_2^- \right\rangle \left( \frac{(\eta-1)\frac{\hbar b}{4\sqrt{\cdots}} + \frac{1}{2}}{E - E_1^{(\pm)}} - \frac{(\eta-1)\frac{\hbar b}{4\sqrt{\cdots}} - \frac{1}{2}}{E - E_2^{(\pm)}} \right)
\end{aligned}
$$

with the four energy poles

$$
\begin{aligned}
E_1^{(\pm)} \ &=\ \mp J_{\mathrm{H}}\hbar^2 - \tfrac{\hbar}{2}(1+\eta)b - \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + \frac{1}{2}(1-\eta)b^2\hbar^2}, \\
E_2^{(\pm)} \ &=\ \mp J_{\mathrm{H}}\hbar^2 - \tfrac{\hbar}{2}(1+\eta)b + \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + \frac{1}{2}(1-\eta)b^2\hbar^2}.
\end{aligned} \tag{A9}
$$

One can apply the spectral theorem (26) to these Green's functions. The corresponding left hand sides are

$$
\begin{array}{lll}
G_{11} \longrightarrow \left\langle S_1^+ S_1^- \right\rangle &=& \frac{\hbar^2}{2} + \hbar \left\langle S_1^z \right\rangle, \\
G_{21} \longrightarrow \left\langle S_1^+ S_2^- \right\rangle, & & \\
\Gamma_{12} \longrightarrow \left\langle S_1^+ S_1^z S_2^- \right\rangle &=& -\frac{\hbar}{2} \left\langle S_1^+ S_2^- \right\rangle, \\
\Gamma_{21} \longrightarrow \left\langle S_2^z S_1^+ S_1^- \right\rangle &=& \frac{\hbar^2}{2} \left\langle S_2^z \right\rangle + \hbar \left\langle S_1^z S_2^z \right\rangle.
\end{array} \tag{A10}
$$

The emerging Bose distribution functions at the right hand side of (26) can be abbreviated:

$$
m_i^{(\pm)} = \frac{1}{\mathrm{e}^{\beta E_i^{(\pm)}} - 1}. \tag{A11}
$$

One ends up with a system of three equations which are linear in the correlation functions we are looking for. The fourth equation is just necessary to calculate $\langle S_2^z \rangle$. It is most instructive to give the corresponding matrix equations for each of the following two special situations separately.

### 1. Ferromagnet ($\eta = +1$)

Here, the expressions for the energy poles are

$$
\begin{array}{llll}
E_1^{(+)} &=& -\hbar b - 2J_{\mathrm{H}}\hbar^2, \quad & E_1^{(-)} \ =\ -\hbar b, \\
E_2^{(+)} &=& -\hbar b, & E_2^{(-)} \ =\ -\hbar b + 2J_{\mathrm{H}}\hbar^2.
\end{array} \tag{A12}
$$

As a consequence $m_1^{(-)} \equiv m_2^{(+)}$. The system of equations is:

$$
\begin{pmatrix} \frac{\hbar^2}{2} \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -1 - \frac{1}{2}m_3 & -\frac{1}{2}\left(m_1^{(+)} - m_2^{(-)}\right) & -\frac{1}{2}\left(m_1^{(+)} - m_2^{(-)}\right) \\ \frac{1}{2}m_4 & \frac{1}{2}\left(m_1^{(+)} - m_2^{(-)}\right) & -1 + \frac{1}{2}\left(m_1^{(+)} - m_2^{(-)}\right) \\ \frac{1}{2}\left(m_1^{(+)} - m_2^{(-)}\right) & \frac{1}{2}m_4 & 1 + \frac{1}{2}m_3 \end{pmatrix} \begin{pmatrix} \hbar \left\langle S_1^z \right\rangle \\ 2 \left\langle S_1^z S_2^z \right\rangle \\ \left\langle S_1^+ S_2^- \right\rangle \end{pmatrix},
$$

where the abbreviations $m_3 = m_1^{(+)} + 2m_1^{(-)} + m_2^{(-)}$ and $m_4 = m_1^{(+)} - 2m_1^{(-)} + m_2^{(-)}$ have been used.

This gives a solution for the correlation functions which can be simplified to the expressions published before by Nolting[15]:

$$\langle S_1^z \rangle = \frac{\hbar}{2} \frac{\exp[\beta\hbar b] - \exp[-\beta\hbar b]}{1 + \exp\left[-2\beta\hbar^2 J_{\mathrm{H}}\right] + \exp[\beta\hbar b] - \exp[-\beta\hbar b]}, \tag{A13}$$

$$\left\langle S_1^+ S_2^- \right\rangle = \frac{\hbar^2}{2} \frac{1 - \exp\left[-2\beta\hbar^2 J_{\mathrm{H}}\right]}{1 + \exp\left[-2\beta\hbar^2 J_{\mathrm{H}}\right] + \exp[\beta\hbar b] - \exp[-\beta\hbar b]}, \tag{A14}$$

$$\langle S_1^z S_2^z \rangle = \frac{\hbar^2}{4} \frac{\exp[\beta\hbar b] - \exp[-\beta\hbar b] - 1 - \exp\left[-2\beta\hbar^2 J_{\mathrm{H}}\right]}{1 + \exp\left[-2\beta\hbar^2 J_{\mathrm{H}}\right] + \exp[\beta\hbar b] - \exp[-\beta\hbar b]}, \tag{A15}$$

$$\langle S_1^z \rangle = \langle S_2^z \rangle \tag{A16}$$

In contrast to his calculations we obtain here the last line as a result and not as an assumption.

### 2. Antiferromagnet ($\eta = -1$)

Here, the expressions for the energy poles are

$$E_1^{(+)} = -J_{\mathrm{H}}\hbar^2 - \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2}, \tag{A17}$$

$$E_1^{(-)} = +J_{\mathrm{H}}\hbar^2 - \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2}, \tag{A18}$$

$$E_2^{(+)} = -J_{\mathrm{H}}\hbar^2 + \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2}, \tag{A19}$$

$$E_2^{(-)} = +J_{\mathrm{H}}\hbar^2 + \sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2} \tag{A20}$$

and consequently

$$\begin{aligned} \Longrightarrow\ & m_1^{(+)} + m_2^{(-)} = -1, \qquad m_1^{(+)} - m_2^{(-)} = 2m_1^{(+)} + 1, \\ & m_2^{(+)} + m_1^{(-)} = -1, \qquad m_2^{(+)} - m_1^{(-)} = 2m_2^{(+)} + 1. \end{aligned}$$

The system of equations is:

$$\begin{pmatrix} \frac{\hbar^2}{2} \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{\hbar b}{\sqrt{\cdots}} \left(m_1^{(+)} - m_2^{(+)}\right) & -\left(m_1^{(+)} + m_2^{(+)} + 1\right) & -\frac{J_{\mathrm{H}}\hbar^2}{\sqrt{\cdots}} \left(m_1^{(+)} - m_2^{(+)}\right) \\ 0 & \frac{J_{\mathrm{H}}\hbar^2}{\sqrt{\cdots}} \left(m_1^{(+)} - m_2^{(+)}\right) & \left(m_1^{(+)} + m_2^{(+)}\right) \\ \frac{J_{\mathrm{H}}\hbar^2}{\sqrt{\cdots}} \left(m_1^{(+)} - m_2^{(+)}\right) & 0 & \frac{\hbar b}{\sqrt{\cdots}} \left(m_1^{(+)} - m_2^{(+)}\right) \end{pmatrix} \begin{pmatrix} \hbar \langle S_1^z \rangle \\ 2 \langle S_1^z S_2^z \rangle \\ \langle S_1^+ S_2^- \rangle \end{pmatrix}$$

Here, the following set correlation functions, which to our knowledge has not been published before, is obtained.

$$\left\langle S_1^+ S_2^- \right\rangle = \frac{J_{\mathrm{H}}\hbar^2}{\sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2}} \cdot \frac{\hbar^2}{2} \cdot \frac{m_1^{(+)} - m_2^{(+)}}{m_1^{(+)} + 4m_1^{(+)}m_2^{(+)} + m_2^{(+)}} \tag{A21}$$

$$\langle S_1^z S_2^z \rangle = -\frac{\hbar^2}{4} \cdot \frac{m_1^{(+)} + m_2^{(+)}}{m_1^{(+)} + 4m_1^{(+)}m_2^{(+)} + m_2^{(+)}} \tag{A22}$$

$$\langle S_1^z \rangle = \frac{b\hbar}{\sqrt{\left(J_{\mathrm{H}}\hbar^2\right)^2 + (b\hbar)^2}} \cdot \frac{\hbar}{2} \cdot \frac{m_1^{(+)} - m_2^{(+)}}{m_1^{(+)} + 4m_1^{(+)}m_2^{(+)} + m_2^{(+)}} \tag{A23}$$

$$\langle S_2^z \rangle = -\langle S_1^z \rangle \tag{A24}$$

The last line, which characterizes an antiferromagnet, is again a result of the calculations.

* Electronic address: hickel@physik.hu-berlin.de


[1] C. Zener, Phys. Rev. **81**, 440 (1951).

[2] D. Meyer, C. Santos and W. Nolting, J. Phys.: Condens. Matter **13**, 2531 (2001).

[3] B.S. Shastry and D.C. Mattis, Phys. Rev. B **24**, 5340 (1981).

[4] W. Nolting, S. Mathi Jaya and S. Rex, Phys. Rev. B **54**, 14 455 (1996).

[5] M. Sigrist, H. Tsunetsugu and K. Ueda, Phys. Rev. Lett. **67**, 2211 (1991).

[6] W. Nolting and M. Matlak, phys. stat. sol. (b) **123**, 155 (1984).

[7] A. Harris and R. Lange, Phys. Rev. **157**, 295 (1967).

[8] M. Matlak, 1999 (private communication).

[9] D. Sénéchal, D. Perez and M. Pioro-Ladrière, Phys. Rev. Lett. **84**, 522 (2000).

[10] A. Avella, F. Mancini, T. Saikawa, cond-mat/0103610 (2001).

[11] R. Schumann, cond-mat/0101476 (2001).

[12] M. Matlak, A. Ramakanth and K. Skrobiś, Z. Phys. B **45**, 331 (1982).

[13] T. Hickel, J. Röseler and W. Nolting, cond-mat/0211670 or:
T. Hickel *Exact Statements on the correlated Kondo-Lattice Model* (diploma thesis, Department of Physics, Humboldt-Universität zu Berlin, 2001).

[14] P.J. Jensen and F. Aguilera-Granja, Phys. Lett. A **269**, 158 (2000).

[15] Wolfgang Nolting, *Quantentheorie des Magnetismus 2* (B. G. Teubner, Stuttgart, 1986).